



TITLE:

COMPOSITE ITERATIVE METHODS FOR A
GENERAL SYSTEM OF VARIATIONAL
INEQUALITIES AND FIXED POINT PROBLEMS
(Study on Nonlinear Analysis and Convex
Analysis)

AUTHOR(S):

Jung, Jong Soo

CITATION:

Jung, Jong Soo. COMPOSITE ITERATIVE METHODS FOR A GENERAL SYSTEM OF VARIATIONAL INEQUALITIES AND FIXED POINT PROBLEMS (Study on Nonlinear Analysis and Convex Analysis). 数理解析研究所講究録 2019, 2112: 13-19

ISSUE DATE:

2019-04

URL:

<http://hdl.handle.net/2433/251984>

RIGHT:

COMPOSITE ITERATIVE METHODS FOR A GENERAL SYSTEM OF VARIATIONAL INEQUALITIES AND FIXED POINT PROBLEMS

JONG SOO JUNG

DEPARTMENT OF MATHEMATICS, DONG-A UNIVERSITY

ABSTRACT. In this paper, we introduce two composite iterative methods (one implicit method and one explicit method) for finding a common element of the solution set of a general system of variational inequalities for continuous monotone mappings and the fixed point set of a continuous pseudocontractive mapping in a Hilbert space. First, this system of variational inequalities is proven to be equivalent to a fixed point problem of nonexpansive mapping. Then we establish strong convergence of the sequence generated by the proposed iterative methods to a common element of the solution set and the fixed point set, which is the unique solution of a certain variational inequality.

1. INTRODUCTION

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and induced norm $\| \cdot \|$. Let C be a nonempty closed convex subset of H and let $S : C \rightarrow C$ be a self-mapping on C . We denote by $Fix(S)$ the set of fixed points of S .

A mapping $F : C \rightarrow H$ is called *monotone* if

$$\langle x - y, Fx - Fy \rangle \geq 0, \quad \forall x, y \in C.$$

and F is called α -*inverse-strongly monotone* (see [5, 11]) if there exists a positive real number α such that

$$\langle x - y, Fx - Fy \rangle \geq \alpha \|Fx - Fy\|^2, \quad \forall x, y \in C.$$

The class of monotone mappings includes the class of α -inverse-strongly monotone mappings.

A mapping $T : C \rightarrow H$ is said to be *pseudocontractive* if

$$\|Tx - Ty\|^2 \leq \|x - y\|^2 + \|(I - T)x - (I - T)y\|^2, \quad \forall x, y \in C.$$

and T is said to be k -*strictly pseudocontractive* if there exists a constant $k \in [0, 1)$ such that

$$\|Tx - Ty\|^2 \leq \|x - y\|^2 + k\|(I - T)x - (I - T)y\|^2, \quad \forall x, y \in C,$$

where I is the identity mapping. Note that the class of k -strictly pseudocontractive mappings includes the class of nonexpansive mappings as a subclass. That is, T is nonexpansive (i.e., $\|Tx - Ty\| \leq \|x - y\|$, $\forall x, y \in C$) if and only if T is 0-strictly pseudocontractive. Clearly, the class of pseudocontractive mappings includes the class of strictly pseudocontractive mappings as a subclass.

Let F be a nonlinear mapping of C into H . The variational inequality problem (VIP) is to find a $x^* \in C$ such that

$$(1.1) \quad \langle Fx^*, x - x^* \rangle \geq 0, \quad \forall x \in C.$$

2010 *Mathematics Subject Classification.* Primary 47J20 Secondary 47H05, 47H09, 47H10, 47J25, 49J40, 49M05, 65J15.

Key words and phrases. Composite iterative method, general system of variational inequalities, continuous monotone mapping, continuous pseudocontractive mapping, ρ -Lipschitzian, η -strongly monotone mapping, variational inequality, strongly positive bounded linear operator, fixed points.

The results presented in this lecture are collected mainly from the work [8] by the author of this report.

We denote the set of solutions of VIP(1.1) by $VI(C, F)$. The variational inequality problem has been extensively studied in the literature; see [3, 5, 7, 10, 11, 14, 15, 17, 19] and the references therein.

In 2008, Ceng *et al.* [2] considered the following general system of variational inequalities:

$$(1.2) \quad \begin{cases} \langle \lambda F_1 y^* + x^* - y^*, x - x^* \rangle \geq 0, & \forall x \in C \\ \langle \nu F_2 x^* + y^* - x^*, x - y^* \rangle \geq 0, & \forall x \in C, \end{cases}$$

where F_1 and F_2 are an α -inverse-strongly monotone mapping and a β -inverse-strongly monotone mapping, respectively; and $\lambda \in (0, 2\alpha)$ and $\nu \in (0, 2\beta)$ are two constants. For finding an element $Fix(S) \cap \Gamma$, where $S : C \rightarrow C$ is a nonexpansive mapping and Γ is the solution set of the problem (1.2), they introduced a relaxed extragradient method ([9]) and proved strong convergence to a common element of $Fix(S) \cap \Gamma$.

In 2016, Alofi *et al.* [1] also considered the problem (1.2) coupled with the fixed point problem, and introduced two composite iterative algorithms (one implicit algorithm and one explicit algorithm) based on Jung's composite iterative method [6] to find an element $Fix(T) \cap \Gamma$, where $T : C \rightarrow C$ is a k -strictly pseudocontractive mapping and Γ is the solution set of the problem (1.2), and showed strong convergence to a common element of $Fix(T) \cap \Gamma$. The following problems arise:

Question 1. Can we extend the class of inverse-strongly monotone mappings in [1, 2] to the more general class of continuous monotone mappings ?

Question 2. Can we extend the class of nonexpansive mappings in [2] or the class of strictly pseudocontractive mappings in [1] to the more general class of pseudocontractive mappings ?

In this paper, in order to give the affirmative answers to the above two questions, we consider a general system of variational inequalities slightly different from the problem (1.2). More precisely, we introduce the following general system of variational inequalities (GSVI) for two continuous monotone mappings F_1 and F_2 of finding $(x^*, y^*) \in C \times C$ such that

$$(1.3) \quad \begin{cases} \langle \lambda F_1 x^* + x^* - y^*, x - x^* \rangle \geq 0, & \forall x \in C \\ \langle \nu F_2 y^* + y^* - x^*, x - y^* \rangle \geq 0, & \forall x \in C, \end{cases}$$

where $\lambda > 0$ and ν are two constants. The solution set of GSVI(1.3) is denoted by Ω . First, we prove that the problem (1.3) is equivalent to a fixed point problem of nonexpansive mapping. Second, by using Jung's composite iterative algorithms [6], we introduce a composite implicit iterative algorithm and a composite explicit iterative algorithm for finding a common element of $\Omega \cap Fix(T)$, where T is a continuous pseudocontractive mapping. Then we establish strong convergence of these two composite iterative algorithms to a common element of $\Omega \cap Fix(T)$, which is the unique solution of a certain variational inequality related to a minimization problem. As a direct consequence, we obtain strong convergence to a common element of $VI(C, F) \cap Fix(T)$, where F is a continuous monotone mapping.

2. PRELIMINARIES AND LEMMAS

Let H be a real Hilbert space and let C be a nonempty closed convex subset of H . We write $x_n \rightharpoonup x$ to indicate that the sequence $\{x_n\}$ converges weakly to x . $x_n \rightarrow x$ implies that $\{x_n\}$ converges strongly to x .

For every point $x \in H$, there exists a unique nearest point in C , denoted by $P_C(x)$, such that

$$\|x - P_C(x)\| \leq \|x - y\|, \quad \forall y \in C.$$

P_C is called the *metric projection* of H onto C . It is well known that $P_C(x)$ is characterized by the property:

$$(2.1) \quad u = P_C(x) \iff \langle x - u, u - y \rangle \geq 0, \quad \forall x \in H, y \in C.$$

In a Hilbert space H , we have

$$(2.2) \quad \|x - y\|^2 = \|x\|^2 + \|y\|^2 - 2\langle x, y \rangle, \quad \forall x, y \in H.$$

We recall that

- (1) an operator A is said to be *strongly positive* on H if there exists a constant $\bar{\gamma} > 0$ such that

$$\langle Ax, x \rangle \geq \bar{\gamma}\|x\|^2, \quad \forall x \in H;$$

- (ii) a mapping $V : C \rightarrow H$ is said to be *l -Lipschitzian* if there exists a constant $l \geq 0$ such that

$$\|Vx - Vy\| \leq l\|x - y\|, \quad \forall x, y \in C;$$

- (iii) a mapping $G : C \rightarrow H$ is said to be *ρ -strongly monotone* if there exists a constant $\rho > 0$ such that

$$\langle Gx - Gy, x - y \rangle \geq \rho\|x - y\|^2, \quad \forall x, y \in C.$$

The following lemma is an immediate consequence of an inner product.

Lemma 2.1. *In a real Hilbert space H , there holds the following inequality*

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, x + y \rangle, \quad \forall x, y \in H.$$

We need the following lemmas for the proof of our main results.

Lemma 2.2 ([16]). *Let $\{s_n\}$ be a sequence of non-negative real numbers satisfying*

$$s_{n+1} \leq (1 - \omega_n)s_n + \omega_n\delta_n + \nu_n, \quad \forall n \geq 1,$$

where $\{\omega_n\}$, $\{\delta_n\}$, and $\{\nu_n\}$ satisfy the following conditions:

- (i) $\{\omega_n\} \subset [0, 1]$ and $\sum_{n=1}^{\infty} \omega_n = \infty$ or, equivalently, $\prod_{n=1}^{\infty} (1 - \omega_n) = 0$;
- (ii) $\limsup_{n \rightarrow \infty} \delta_n \leq 0$ or $\sum_{n=1}^{\infty} \omega_n |\delta_n| < \infty$;
- (iii) $\nu_n \geq 0$ ($n \geq 1$), $\sum_{n=1}^{\infty} \nu_n < \infty$.

Then $\lim_{n \rightarrow \infty} s_n = 0$.

Lemma 2.3 ([4]). (Demiclosedness principle) *Let C be a nonempty closed convex subset of a real Hilbert space H , and let $S : C \rightarrow C$ be a nonexpansive mapping. Then, the mapping $I - S$ is demiclosed. That is, if $\{x_n\}$ is a sequence in C such that $x_n \rightharpoonup x^*$ and $(I - S)x_n \rightarrow y$, then $(I - S)x^* = y$.*

Lemma 2.4 ([12]). *Let H be a real Hilbert space. Let $A : H \rightarrow H$ be a strongly positive bounded linear operator with a constant $\bar{\gamma} > 1$. Then*

$$\langle (A - I)x - (A - I)y, x - y \rangle \geq (\bar{\gamma} - 1)\|x - y\|^2, \quad \forall x, y \in C.$$

That is, $A - I$ is strongly monotone with a constant $\bar{\gamma} - 1$.

Lemma 2.5 ([12]). *Assume that A is a strongly positive bounded linear operator on H with a coefficient $\bar{\gamma} > 0$ and $0 < \zeta \leq \|A\|^{-1}$. Then $\|I - \zeta A\| \leq 1 - \zeta\bar{\gamma}$.*

Lemma 2.6 ([17]). *Let H be a real Hilbert space. Let $G : H \rightarrow H$ be a ρ -Lipschitzian and η -strongly monotone mapping with constants $\rho, \eta > 0$. Let $0 < \mu < \frac{2\eta}{\rho^2}$ and $0 < t < \sigma \leq 1$. Then $S := \sigma I - t\mu G : H \rightarrow H$ is a contractive mapping with constant $\sigma - t\tau$, where $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\rho^2)}$.*

The following lemmas are Lemma 2.3 and Lemma 2.4 of Zegeye [18], respectively.

Lemma 2.7 ([18]). *Let C be a closed convex subset of a real Hilbert space H . Let $F : C \rightarrow H$ be a continuous monotone mapping. Then, for $r > 0$ and $x \in H$, there exists $z \in C$ such that*

$$\langle y - z, Fz \rangle + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \quad \forall y \in C.$$

For $r > 0$ and $x \in H$, define $F_r : H \rightarrow C$ by

$$F_r x = \left\{ z \in C : \langle y - z, Fz \rangle + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \quad \forall y \in C \right\}.$$

Then the following hold:

- (i) F_r is single-valued;
- (ii) F_r is firmly nonexpansive, that is,

$$\|F_r x - F_r y\|^2 \leq \langle x - y, F_r x - F_r y \rangle, \quad \forall x, y \in H;$$
- (iii) $\text{Fix}(F_r) = \text{VI}(C, F)$;
- (iv) $\text{VI}(C, F)$ is a closed convex subset of C .

Lemma 2.8 ([18]). *Let C be a closed convex subset of a real Hilbert space H . Let $T : C \rightarrow H$ be a continuous pseudocontractive mapping. Then, for $r > 0$ and $x \in H$, there exists $z \in C$ such that*

$$\langle y - z, Tz \rangle - \frac{1}{r} \langle y - z, (1 + r)z - x \rangle \leq 0, \quad \forall y \in C.$$

For $r > 0$ and $x \in H$, define $T_r : H \rightarrow C$ by

$$T_r x = \left\{ z \in C : \langle y - z, Tz \rangle - \frac{1}{r} \langle y - z, (1 + r)z - x \rangle \leq 0, \quad \forall y \in C \right\}.$$

Then the following hold:

- (i) T_r is single-valued;
- (ii) T_r is firmly nonexpansive, that is,

$$\|T_r x - T_r y\|^2 \leq \langle x - y, T_r x - T_r y \rangle, \quad \forall x, y \in H;$$
- (iii) $\text{Fix}(T_r) = \text{Fix}(T)$;
- (iv) $\text{Fix}(T)$ is a closed convex subset of C .

3. MAIN RESULTS

Throughout the rest of this paper, we always assume the following:

- H is a real Hilbert space;
- C is a nonempty closed subspace subset of H ;
- $A : C \rightarrow C$ is a strongly positive linear bounded self-adjoint operator with a constant $\bar{\gamma} \in (1, 2)$;
- $V : C \rightarrow C$ is l -Lipschitzian with constant $l \in [0, \infty)$;
- $G : C \rightarrow C$ is a ρ -Lipschitzian and η -strongly monotone mapping with constants $\rho > 0$ and $\eta > 0$;
- Constants μ , l , τ , and γ satisfy $0 < \mu < \frac{2\eta}{\rho^2}$ and $0 \leq \gamma l < \tau$, where $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\rho^2)}$;
- F_1 and $F_2 : C \rightarrow H$ are continuous monotone mapping;
- Ω is the solution set of GSVI (1.3) for F_1 and F_2 ;
- $F_{1\lambda} : H \rightarrow C$ is a mapping defined by

$$F_{1\lambda} x = \left\{ z \in C : \langle y - z, F_1 z \rangle + \frac{1}{\lambda} \langle y - z, z - x \rangle \geq 0, \quad \forall y \in C \right\}$$

for $\lambda > 0$;

- $F_{2\nu} : H \rightarrow C$ is a mapping defined by

$$F_{2\nu}x = \left\{ z \in C : \langle y - z, F_2z \rangle + \frac{1}{\nu} \langle y - z, z - x \rangle \geq 0, \quad \forall y \in C \right\}$$

for $\nu > 0$;

- $R : H \rightarrow C$ is a mapping defined by $Rx = F_{1\lambda}F_{2\nu}x$ for each $x \in H$;
- $T : C \rightarrow C$ is a continuous pseudocontractive mapping such that $Fix(T) \neq \emptyset$;
- $T_{r_t} : H \rightarrow C$ is a mapping defined by

$$T_{r_t}x = \left\{ z \in C : \langle y - z, Tz \rangle - \frac{1}{r_t} \langle y - z, (1 + r_t)z - x \rangle \leq 0, \quad \forall y \in C \right\}$$

for $r_t \in (0, \infty)$, $t \in (0, 1)$, and $\liminf_{t \rightarrow 0} r_t > 0$;

- $T_{r_n} : H \rightarrow C$ is a mapping defined by

$$T_{r_n}x = \left\{ z \in C : \langle y - z, Tz \rangle - \frac{1}{r_n} \langle y - z, (1 + r_n)z - x \rangle \leq 0, \quad \forall y \in C \right\}$$

for $r_n \in (0, \infty)$ and $\liminf_{n \rightarrow \infty} r_n > 0$;

- $\Omega \cap Fix(T) \neq \emptyset$.

By Lemma 2.7 and Lemma 2.8, we note that $F_{1\lambda}$, $F_{2\nu}$, T_{r_t} , and T_{r_n} are nonexpansive, and $Fix(T_{r_n}) = Fix(T) = Fix(T_{r_t})$.

First, we prove that the problem (1.3) is equivalent to a fixed point problem of nonexpansive mapping.

Proposition 3.1. *Let C be a closed convex subset of a real Hilbert space H . For given $x^*, y^* \in C$, (x^*, y^*) is a solution of GSVI(1.3) for continuous monotone mappings F_1 and F_2 if and only if x^* is a fixed point of the mapping $R : H \rightarrow C$ defined by*

$$Rx = F_{1\lambda}F_{2\nu}x, \quad \forall x \in H,$$

where $y^* = F_{2\nu}x^*$.

First, we introduce the following composite algorithm that generates a net $\{x_t\}$ in an implicit way:

$$(3.1) \quad x_t = (I - \theta_t A)T_{r_t}Rx_t + \theta_t[t\gamma Vx_t + (I - t\mu G)T_{r_t}Rx_t],$$

where $t \in (0, \min\{1, \frac{2-\bar{\gamma}}{\tau-\bar{\gamma}l}\})$ and $\theta_t \in (0, \|A\|^{-1}]$.

We summarize the basic properties of $\{x_t\}$, which can be proved by the same method as in [6].

Proposition 3.2. *Let $\{x_t\}$ be defined via (3.1). Then*

- (i) $\{x_t\}$ is bounded for $t \in (0, \min\{1, \frac{2-\bar{\gamma}}{\tau-\bar{\gamma}l}\})$;
- (ii) $\lim_{t \rightarrow 0} \|x_t - T_{r_t}Rx_t\| = 0$ provided $\lim_{t \rightarrow 0} \theta_t = 0$;
- (iii) $\lim_{t \rightarrow 0} \|x_t - y_t\| = 0$, where $y_t = t\gamma Vx_t + (I - t\mu G)T_{r_t}Rx_t$;
- (iv) $\lim_{t \rightarrow 0} \|x_t - Rx_t\| = 0$;
- (v) x_t defines a continuous path from $(0, \min\{1, \frac{2-\bar{\gamma}}{\tau-\bar{\gamma}l}\})$ into H provided $\theta_t : (0, \min\{1, \frac{2-\bar{\gamma}}{\tau-\bar{\gamma}l}\}) \rightarrow (0, \|A\|^{-1}]$ is continuous, and $r_t : (0, \min\{1, \frac{2-\bar{\gamma}}{\tau-\bar{\gamma}l}\}) \rightarrow (0, \infty)$ is continuous.

We obtain the following theorem for strong convergence of the net $\{x_t\}$ as $t \rightarrow 0$, which guarantees the existence of solutions of the variational inequality (3.2) below.

Theorem 3.3. *Let the net $\{x_t\}$ be defined via (3.1). If $\lim_{t \rightarrow 0} \theta_t = 0$, then x_t converges strongly to \tilde{x} in $\Omega \cap \text{Fix}(T)$ as $t \rightarrow 0$, which solves the variational inequality*

$$(3.2) \quad \langle (A - I)\tilde{x}, \tilde{x} - p \rangle \leq 0, \quad \forall p \in \Omega \cap \text{Fix}(T),$$

Equivalently, we have

$$P_{\Omega \cap \text{Fix}(T)}(2I - A)\tilde{x} = \tilde{x}.$$

Now, we propose the following composite algorithm which generates a sequence in an explicit way:

$$(3.3) \quad \begin{cases} y_n = \alpha_n \gamma V x_n + (I - \alpha_n \mu G) T_{r_n} R x_n, \\ x_{n+1} = (I - \beta_n A) T_{r_n} R x_n + \beta_n y_n, \quad \forall n \geq 0, \end{cases}$$

where $\{\alpha_n\} \in [0, 1]$; $\{\beta_n\} \subset (0, 1]$; $\{r_n\} \subset (0, \infty)$; and $x_0 \in C$ is an arbitrary initial guess, and establish strong convergence of this sequence to $\tilde{x} \in \Omega \cap \text{Fix}(T)$, which is the unique solution of the variational inequality (3.2).

Theorem 3.4. *Let $\{x_n\}$ be the sequence generated by the explicit algorithm (3.3). Let $\{\alpha_n\}$, $\{\beta_n\}$, and $\{r_n\}$ satisfy the following conditions:*

- (C1) $\{\alpha_n\} \subset [0, 1]$ and $\{\beta_n\} \subset (0, 1]$, $\alpha_n \rightarrow 0$ and $\beta_n \rightarrow 0$ as $n \rightarrow \infty$;
- (C2) $\sum_{n=0}^{\infty} \beta_n = \infty$;
- (C3) $\sum_{n=0}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty$, and $|\beta_{n+1} - \beta_n| \leq o(\beta_{n+1}) + \sigma_n$, $\sum_{n=0}^{\infty} \sigma_n < \infty$ (the perturbed control condition);
- (C4) $\{r_n\} \subset (0, \infty)$, $\liminf_{n \rightarrow \infty} r_n > 0$, and $\sum_{n=0}^{\infty} |r_{n+1} - r_n| < \infty$.

Then $\{x_n\}$ converges strongly to $\tilde{x} \in \Omega \cap \text{Fix}(T)$, which is the unique solution of the variational inequality (3.2).

Taking $G \equiv I$, $\mu = 1$, and $\gamma = 1$ in Theorem 3.5, we obtain the following corollary.

Corollary 3.5. *Let $\{x_n\}$ be generated by the following iterative algorithm:*

$$\begin{cases} y_n = \alpha_n V x_n + (1 - \alpha_n) T_{r_n} R x_n, \\ x_{n+1} = (I - \beta_n A) T_{r_n} R x_n + \beta_n y_n, \quad \forall n \geq 0. \end{cases}$$

Assume that the sequences $\{\alpha_n\}$, $\{\beta_n\}$, and $\{r_n\}$ satisfy the conditions (C1) – (C4) in Theorem 3.5. Then $\{x_n\}$ converges strongly to $\tilde{x} \in \Omega \cap \text{Fix}(T)$, which is the unique solution of the variational inequality (3.2).

Remark 3.6. 1) The $\tilde{x} \in \Omega \cap \text{Fix}(T)$ in our results is the unique solution of minimization problem

$$(3.4) \quad \min_{x \in D} \frac{1}{2} \langle (A - I)x, x \rangle,$$

where the constraint set D is $\Omega \cap \text{Fix}(T)$. In fact, the variational inequality (3.2) is the optimality condition for the minimization problem (3.4). Thus, for finding an element of $\Omega \cap \text{Fix}(T)$, where T is a continuous pseudocontractive mapping, and F_1 and F_2 are continuous monotone mappings, Theorem 3.4, Theorem 3.5 and Corollary 3.6 are new ones different from previous those introduced by some authors (for example, see [1, 2]).

2) Taking $F_1 = F_2 = F$, $\lambda = \nu$ and $x^* = y^*$ in GSVI(1.3) and replacing F_λ by F_{r_n} along with the condition (C4) on $\{r_n\}$, we can obtain a new result, which improves, supplements and develops the corresponding results of [3, 5, 7, 14, 15, 19].

ACKNOWLEDGMENT

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2018R1D1A1B07045718).

REFERENCES

- [1] A. S. M. Alofi, A. Latif, A. E. Al-Marzooei, J. C. Yao, *Composite viscosity iterative methods for general systems of variational inequalities and fixed point problem in Hilbert spaces*, J. Nonlinear Convex Anal. **16** (2016), no. 4, 669–682.
- [2] L. C. Ceng, C. Y. Wang, J. C. Yao, *Strong convergence theorems by a relaxed extragradient method for a general system of variational inequalities*, Math. Meth. Oper. Res. **67** (2008), 375–390.
- [3] J. Chen, L. Zhang, T. Fan, *Viscosity approximation methods for nonexpansive mappings and monotone mappings*, J. Math. Anal. Appl. **334** (2007), 1450–1461.
- [4] K. Goebel, W. A. Kirk, *Topics in Metric Fixed Point Theory*, in: Cambridge Stud. Adv. Math., vol 28, Cambridge Univ. Press, 1990.
- [5] H. Iiduka, W. Takahashi, *Strong convergence theorems for nonexpansive mappings and inverse-strongly monotone mappings*, Nonlinear Anal. **61** (2005), 341–350.
- [6] J. S. Jung, *A general composite iterative method for strictly pseudocontractive mappings in Hilbert spaces*, Fixed Point Theory Appl. **2014** (2014), doi:10.1186/1687-1812-2014-173.
- [7] J. S. Jung, *A composite extragradient-like algorithm for inverse-strongly monotone mappings and strictly pseudocontractive mappings*, Linear and Nonlinear Anal. **1** (2015), no. 2, 271–285.
- [8] J. S. Jung, *Strong convergence of some iterative algorithms for a general system of variational inequalities* J. Nonlinear Sci. Appl. **10** (2017), 3887–3902.
- [9] G. M. Korpelevich, *An extragradient method for finding saddle points and for other problems*, Ekon. Mate. Metody, **12** (1976), 747–756.
- [10] P. L. Lions, G. Stampacchia, *Variational inequalities*, Comm. Pure Appl. Math. **20** (1967), 493–517.
- [11] F. Liu, M. Z. Nashed, *Regularization of nonlinear ill-posed variational inequalities and convergence rates*, Set-Valued Anal. **6** (1998), 313–344.
- [12] G. Marino, H. K. Xu, *A general iterative method for nonexpansive mappings in Hilbert spaces*, J. Math. Anal. Appl. **318** (2006), 43–52.
- [13] G. J. Minty, *On the generalization of a direct method of the calculus of variations*, Bull. Amer. Math. Soc. **73** (1967), 315–321.
- [14] W. Takahashi, M. Toyoda, *Weak convergence theorems for nonexpansive mappings and monotone mappings*, J. Optim. Theory Appl. **118** (2003), no. 2, 417–428.
- [15] Y. Tang, *Strong convergence of viscosity approximation methods for the fixed-point of pseudo-contractive and monotone mappings*, Fixed Point Theory Appl. **2013** (2003), 2013:273.
- [16] H. K. Xu, *An iterative algorithm for nonlinear operator*, J. London Math. Soc. **66** (2002), 240–256.
- [17] I. Yamada, *The hybrid steepest descent method for the variational inequality of the intersection of fixed point sets of nonexpansive mappings*, in D. Butnariu, Y. Censor, S. Reich (Eds), Inherently Parallel Algorithm for Feasibility and Optimization, and Their Applications, Kluwer Academic Publishers, Dordrecht, Holland, pp. 473–504, 2001.
- [18] H. Zegeye, *An iterative approximation method for a common fixed point of two pseudocontractive mappings*, International Scholarly Reserach Network ISRN Math. Anal. **2011** (2011) Article ID 621901, 14 pages.
- [19] H. Zegeye, N. Shahzad, *Strong convergence of an iterative method for pseudo-contractive and monotone mappings*, J. Glob. Optim, **54** (2012), 173–184.

DEPARTMENT OF MATHEMATICS, DONG-A UNIVERSITY, BUSAN 49315, KOREA
 E-mail address: jungjs@dau.ac.kr